

AN ELEMENTARY PROOF OF A SERIES EVALUATION IN TERMS OF HARMONIC NUMBERS

HELMUT PRODINGER

For positive integers j , consider

$$S(j) = \sum_{n \geq 1} \frac{1}{2^{2n-1}(2n-1)} \sum_k \binom{2n-1}{k} \frac{1}{k-j-n+\frac{1}{2}}.$$

This quantity arose in [4] and was subsequently evaluated in [3]. Further proofs of the final formula (1) were given in [2, 1].

Here, we give an extremely short and simple proof.

For our analysis, it is better to replace the inner summation index k by $2n-1-k$, and consider

$$T(j) = \sum_{n \geq 1} \frac{1}{2^{2n-1}(2n-1)} \sum_k \binom{2n-1}{k} \frac{1}{k+j-n+\frac{1}{2}},$$

then $S(j) = -T(j)$.

We start from the obvious formula

$$\int_0^1 t^{x-1}(1+t)^{2n-1} dt = \sum_k \binom{2n-1}{k} \frac{1}{k+x},$$

substitute $x = j - n + \frac{1}{2}$ and sum:

$$\begin{aligned} T(j) &= \int_0^1 t^{j-\frac{1}{2}}(1+t)^{-1} \sum_{n \geq 1} \frac{1}{2^{2n-1}(2n-1)} t^{-n}(1+t)^{2n} dt \\ &= \int_0^1 t^{j-\frac{1}{2}}(1+t)^{-1} \frac{1+t}{\sqrt{t}} \log \frac{1+\sqrt{t}}{1-\sqrt{t}} dt \\ &= 2 \int_0^1 w^{2j-1} \log \frac{1+w}{1-w} dw \\ &= 4 \int_0^1 w^{2j-1} \sum_{k \geq 1} \frac{w^{2k-1}}{2k-1} dw \\ &= 4 \sum_{k \geq 1} \frac{1}{(2k-1)(2k+2j-1)} \\ &= \frac{2}{j} \sum_{k \geq 1} \left(\frac{1}{2k-1} - \frac{1}{2k+2j-1} \right) \\ &= \frac{2}{j} \sum_{k=1}^j \frac{1}{2k-1}. \end{aligned}$$

Hence

$$S(j) = -\frac{2}{j} \sum_{k=1}^j \frac{1}{2k-1}. \quad (1)$$

It is not hard to show that it is allowed to interchange integration and summation.

REFERENCES

- [1] W. Chu and L. De Donno. Transformation on infinite double series and applications to harmonic number identities. *Appl. Algebra Engrg. Comm. Comput.*, 15:339–348, 2005.
- [2] C. Krattenthaler. Hypergeometric proof of a curious identity of Lyons, Paule and Riese. *South East Asian J. Math. Math. Sci.*, 3:63–67, 2004.
- [3] R. Lyons, P. Paule, and A. Riese. A computer proof of a series evaluation in terms of harmonic numbers. *Appl. Algebra Engrg. Comm. Comput.*, 13:327–333, 2002.
- [4] R. Lyons and J. E. Steif. Stationary determinantal processes: phase multiplicity, Bernoullicity, entropy, and domination. *Duke Math. J.*, 120:515–575, 2003.

HELMUT PRODINGER, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF STELLENBOSCH, 7602 STELLENBOSCH, SOUTH AFRICA

E-mail address: `hproding@sun.ac.za`